

DevelopingAveragesStarters: key vocabulary

average	a single value chosen to stand for a whole set of data
bimodal	having two values that appear most often in a list of numbers.
common denominator	a bottom number shared by two or more fractions, used for adding or comparing them.
data	the set of values collected
even	able to be divided exactly by 2.
fraction	a number written as one whole split into equal parts, with a number of those parts taken
frequency	how many times something happens
improper fraction ..	a fraction with a top number larger than or equal to its bottom number
integer	a whole number, which may be positive, negative or zero
mean	the average found by adding all the values and dividing by how many there are
median	the middle value when the data is put in order
mode	the value that appears most often
numerator	the number above the line in a fraction, showing how many equal parts are taken
odd	a whole number that cannot be split exactly into two equal whole numbers.
outlier	a value far away from all the others
positive	a number greater than zero.
range	the difference between the largest and smallest values
set	a collection of numbers or objects grouped together.
symmetric	balanced so that one side is a mirror image of the other.
unique	being the only one; here, no other value is tied with the mode.
value	the number that a letter or symbol stands for

Match the word to its meaning

average	having two values that appear most often in a list of numbers.
bimodal	the set of values collected
common denominator	how many times something happens
data	able to be divided exactly by 2.
even	the average found by adding all the values and dividing by how many there are
fraction	a single value chosen to stand for a whole set of data
frequency	a number written as one whole split into equal parts, with a number of those parts taken
improper fraction	a bottom number shared by two or more fractions, used for adding or comparing them.
integer	a whole number that cannot be split exactly into two equal whole numbers.
mean	a whole number, which may be positive, negative or zero
median	the number above the line in a fraction, showing how many equal parts are taken
mode	a fraction with a top number larger than or equal to its bottom number
numerator	a number greater than zero.
odd	the difference between the largest and smallest values
outlier	the middle value when the data is put in order
positive	balanced so that one side is a mirror image of the other.
range	the value that appears most often
set	the number that a letter or symbol stands for
symmetric	a collection of numbers or objects grouped together.
unique	a value far away from all the others
value	being the only one; here, no other value is tied with the mode.

Match the word to its meaning: answers

average	having two values that appear most often in a list of numbers.
bimodal	the set of values collected
common denominator	how many times something happens
data	able to be divided exactly by 2.
even	the average found by adding all the values and dividing by how many there are
fraction	a single value chosen to stand for a whole set of data
frequency	a number written as one whole split into equal parts, with a number of those parts taken
improper fraction	a bottom number shared by two or more fractions, used for adding or comparing them.
integer	a whole number that cannot be split exactly into two equal whole numbers.
mean	a whole number, which may be positive, negative or zero
median	the number above the line in a fraction, showing how many equal parts are taken
mode	a fraction with a top number larger than or equal to its bottom number
numerator	a number greater than zero.
odd	the difference between the largest and smallest values
outlier	the middle value when the data is put in order
positive	balanced so that one side is a mirror image of the other.
range	the value that appears most often
set	the number that a letter or symbol stands for
symmetric	a collection of numbers or objects grouped together.
unique	a value far away from all the others
value	being the only one; here, no other value is tied with the mode.